

Mammalia, Didelphimorphia and Rodentia, central Santa Fe Province, Argentina

Pablo Teta^{1*} and Ulyses F. J. Pardiñas²

1 Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Avenida Ángel Gallardo 470, (C1405DJR) Buenos Aires, Argentina.

2 Unidad de Investigación Diversidad, Sistemática y Evolución, Centro Nacional Patagónico, CC 128, 9120 Puerto Madryn, Chubut, Argentina.

* Corresponding author. E-mail: anthea@yahoo.com.ar

ABSTRACT: Three owl pellets samples collected in the localities of Pedro Gómez Cello (= Estación Km. 197; 30°02'14" S, 60°18'56" W), Colonia Silva (= Estación Abipones; 30°26'59" S, 60°25'58" W) and Jacinto L. Arauz (30°44'01" S, 60°58'31" W), Province of Santa Fe, Argentina, were studied. We registered 11 genera of small mammals, including Didelphidae marsupials (1 species), and Caviidae (1), Cricetidae (10), and Muridae (1) rodents. We documented the southernmost record for *Pseudoryzomys simplex* and the first and second recording localities for Santa Fe of *Oligoryzomys nigripes* and *Graomys chacoensis*, respectively.

INTRODUCTION

The Great American Chaco occupies more than 1,000,000 km² in the political territories of Argentina, Brazil, Bolivia and Paraguay (Olson *et al.* 2001). This eco-region is divided in an occidental dry sector and in an eastern humid one, which reaches the north-central part of Santa Fe Province (Argentina), being its southernmost expression (Burkart *et al.* 1999). At this place, the Humid Chaco is intermixed with the floristic communities of the Espinal, an endemic eco-region of Argentina (Burkart *et al.* 1999). Small mammal assemblages for this sector, except for some minor references (*e.g.* Massoia *et al.* 1995; Pardiñas *et al.* 2004; Pautasso 2008), remain almost unknown (Galliari and Goin 1993). This situation is better reflected in the scarce number of recording localities for marsupials and rodents with voucher specimens in the central and northern parts of the Santa Fe Province (see Pautasso 2008). In this note, we documented the small terrestrial mammal species of three localities placed in the transitional area between the Humid Chaco and the Espinal. In addition, we make some biogeographical comments regarding the reported species.

MATERIALS AND METHODS

We studied skull and mandible remains found in three Barn Owl (*Tyto alba*; Aves, Tytonidae) pellet samples collected in the localities of: 1- Pedro Gómez Cello (= Estación Km. 197; 30°02'14" S, 60°18'56" W), 2- Colonia Silva (= Estación Abipones; 30°26'59" S, 60°25'58" W) and 3- Jacinto L. Arauz (30°44'01" S, 60°58'31" W) (Figure 1). Taxonomic identifications were made through the comparison with voucher specimens housed in the *Colección de Mamíferos del Centro Nacional Patagónico* (Puerto Madryn, Chubut, Argentina) and literature. Fragmentary remains of small specimens of *Calomys* and *Oligoryzomys*, and young individuals of *Rattus* are difficult to identify to species level (*e.g.*, Massoia 1973; Pardiñas and Lezcano 1995), thus we used an open taxonomy. Studied

materials are deposited in the *Colección de Material de Egagrópilas y Afines "Elio Massoia"* of the *Centro Nacional Patagónico*, under the access numbers CNP-E 196 (Colonia Silva), CNP-E 200 (Pedro Gómez Cello) and CNP-E 460 (Jacinto L. Arauz).




FIGURE 1. Map of the study area, central-eastern Santa Fe Province, Argentina: 1. Pedro Gómez Cello; 2. Colonia Silva; 3. Jacinto L. Arauz.

RESULTS AND DISCUSSION

Micromammal assemblages for the three studied samples are given in Table 1. The sigmodontine rodent *Akodon azarae* (Fischer, 1829) was the best represented species, followed by moderate frequencies of *Calomys* spp.,

a small form of the genus *Oligoryzomys*, *Holochilus chacarius* Thomas, 1906, and *Necomys lasiurus* (Lund, 1840) (Table 1). Both *A. azarae* and *Calomys* spp. [including *C. laucha* (Fischer, 1814) and *C. musculinus* (Thomas, 1913)] are abundant species in the grasslands, pasturelands, and agro ecosystems of the Chaco-Pampean plains (e.g. Kravetz *et al.* 1986). The record in sympatry between *Holochilus brasiliensis* (Desmarest, 1819) and *H. chacarius* enlarges to the west the co-occurrence area for these species in Argentina, that is extended like a fringe in both sides of the Paraná River from Eastern Formosa and northwestern Corrientes (Pardiñas and Teta 2005; Pardiñas *et al.* 2005) to the north of the Buenos Aires (Voglino *et al.* 2005). Judging by the studied samples and literature, *N. lasiurus*, including *N. benefactus* (Thomas, 1919) and *N. temchuki* (Massoia, 1982) (see D’Elía *et al.* 2008), has a more or less continuous distribution with moderately abundant populations in northwestern Argentina (e.g. Pardiñas and Teta 2005), that are gradually fragmented towards the Pampean and Espinal eco-regions (where its participation in the communities is much less conspicuous; see Galliari and Pardiñas 2000). Two small [*O. flavescens* (Waterhouse, 1837) and *O. fornesi* (Massoia, 1973)] and morphologically similar species of the rodent genus *Oligoryzomys* were documented for Santa Fe Province (Massoia *et al.* 1995; Romano *et al.* 2002). Its distributional ranges, especially in middle latitudes of the provinces, are far from being well established. *Oligoryzomys nigripes* (Olfers, 1818) is found at the first time in the Province of Santa Fe; the records for Pedro Gómez Cello and Jacinto L. Arauz enlarges ca. de 250 km to the SW the known distribution of this rodent. In the case of *Graomys chacoensis* (Allen, 1901), the only previous record for the province corresponds to the Departamento La Capital [cited as *G. griseoflavus* by Pautasso (2008)], being the record presented here the second for the province (cf. Ferro and Martinez 2009). This finding also enlarges the known distributional range of this sigmodontine more

TABLE 1. Terrestrial micromammals found in Barn Owl pellet samples in the localities of Pedro Gómez Cello, Colonia Silva, and Jacinto L. Arauz (Santa Fe Province, Argentina).

	Pedro Gómez Cello	Colonia Silva	Jacinto L. Arauz
<i>Akodon azarae</i>	53	71	61
<i>Calomys</i> cf. <i>C. laucha</i> - <i>C. musculinus</i>	8	25	68
<i>Cavia aperea</i>	8	-	1
<i>Cryptonanus chacoensis</i>	8	-	-
<i>Graomys chacoensis</i>	-	-	3
<i>Holochilus brasiliensis</i>	-	-	3
<i>Holochilus chacarius</i>	35	11	4
<i>Necomys lasiurus</i>	11	16	14
<i>Oligoryzomys</i> cf. <i>O. flavescens</i> - <i>O. fornesi</i>	25	27	21
<i>Oligoryzomys nigripes</i>	5	-	7
<i>Pseudoryzomys simplex</i>	25	-	-
<i>Rattus</i> sp.	1	-	-
<i>Scapteromys aquaticus</i>	6	-	-
	185	150	182

than 300 km to the east. The record for *Pseudoryzomys simplex* (Winge, 1887) in Pedro Gómez Cello constitutes the southernmost reference for this species (Pardiñas *et al.* 2004). Finally, in the case of *Scapteromys aquaticus* Thomas, 1920 the studied materials from Pedro Gómez Cello extend its distribution ca. 45 km to the west.

Within the order Didelphimorphia, the unique recorded species was *Cryptonanus chacoensis* (Tate, 1931) in Pedro Gómez Cello. In Santa Fe Province, this small didelphid had only two previous records in the Departamento La Capital, near the coast of the Paraná River (Pautasso 2008).

Part of the records documented in this note enlarges significantly the known distributional range of some sigmodontine rodent species. In the other cases they contribute with a better delimitation of the distributional ranges of species that have a few adequately documented records (e.g. *C. chacoensis*). Some questions posed in this note (e.g. how many small species of *Oligoryzomys* are present in Santa Fe? What are their distributions?) emphasize the need of intensive fieldworks in Santa Fe province. This is especially important because this region constitutes the southernmost distributional limits of some Brazilian small mammals (e.g. *O. fornesi*, *P. simplex*) within a rich biogeographical scenario produced by the intermingled of two main eco-regions, the Chaco and the Espinal.

ACKNOWLEDGMENTS: C. Galliari, D. Podestá, F. Cremonte, J. Notarnicola and S. Cirignoli helped us during field work conducted in Santa Fe and in some cases assisted during laboratory tasks. L. Antoniazzi collected and kindly offered the sample from Jacinto L. Arauz. Two anonymous reviewers made valuable comments on a first draft of this manuscript. Economic support for this research was provided by CONICET.

LITERATURE CITED

Burkart, R., N.O. Bárbaro, R.O. Sánchez and D.A. Gómez. 1999. *Eco-regiones de la Argentina*. Buenos Aires: Instituto Nacional de Tecnología Agropecuaria, PRODIA. 42 p.

D’Elía, G., U.F.J. Pardiñas, J.P. Jayat and J. Salazar-Bravo. 2008. Systematics of *Necomys* (Rodentia, Cricetidae, Sigmodontinae): species limits and groups, with comments on historical biogeography. *Journal of Mammalogy* 89(3): 778-790.

Ferro, L.I., and J.J. Martínez. 2009. Molecular and morphometrics evidence validates a Chacoan species of the grey-eared mice genus *Graomys* (Rodentia: Cricetidae: Sigmodontinae). *Mammalia* 73(3): 265-271.

Galliari, C.A. and F. Goin. 1993. Conservación de la Biodiversidad en la Argentina: el caso de los mamíferos; p. 367-400. In F. Goin and R. Goñi (ed.). *Elementos de Política Ambiental*. La Plata: Honorable Cámara de Diputados de la provincia de Buenos Aires.

Galliari, C.A. and U.F.J. Pardiñas. 2000. Taxonomy and distribution of the sigmodontine rodents of genus *Necomys* in central Argentina and Uruguay. *Acta Theriologica* 45(2): 211-232.

Kravetz, F.O., R.E. Percich, G.A. Zuleta, M.A. Callelo and M.C. Weissembacher. 1986. Distribution of Junin Virus and its reservoirs. A tool for Argentine Hemorrhagic Fever risk. Evaluation in non-endemic areas. *Interiencia* 11(4): 185-188.

Massoia, E., G. Aprile, and B. Lartigau. 1995. Análisis de regurgitados de *Tyto alba* de Estación Santa Margarita, departamento 9 de Julio, provincia de Santa Fe. *Boletín Científico de la asociación para la Protección de la Naturaleza* 9(27): 19-21.

Olson, D.M., E. Dinerstein, E.D. Wikramanayake, N.D. Burgués, G.V.N. Powell, E.C. Underwood, J.A. D’Amico, H.E. Strand, J.C. Morrison, C.J. Loucks, T.F. Allnutt, J.F. Lamoreux, T.H. Ricketts, I. Itoua, W.W. Wettengel, Y. Kura and H. Kassem. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *BioScience* 51(11): 933-938.

Pardiñas, U.F.J. and M. Lezcano. 1995. Cricétidos (Mammalia: Rodentia) del Pleistoceno tardío del nordeste de la provincia de Buenos Aires (Argentina): aspectos sistemáticos y paleoambientales. *Ameghiniana* 32(3): 249-265.

Pardiñas, U.F.J., C.A. Galliari and S. Cirignoli. 2004. Distribution of *Pseudoryzomys simplex* (Rodentia: Cricetidae) in Argentina. *Mastozoología Neotropical* 11(1): 105-108.

- Pardiñas, U.F.J. and P. Teta. 2005. Roedores sigmodontinos del Chaco Húmedo de Formosa, Argentina: aspectos taxonómicos y distribución geográfica; p. 501-517. *In* Di Giacomo, A.G. and S.F. Krapovickas (ed.). *Historia Natural y Paisaje de la Reserva Natural El Bagual, provincia de Formosa, Argentina. Inventario de la fauna de vertebrados y de la flora vascular de un área protegida del Chaco húmedo. Temas de Naturaleza y Conservación 4*. Buenos Aires: Aves Argentinas / Asociación Ornitológica del Plata.
- Pardiñas, U.F.J., P. Teta and S. Heinonen Fortabat. 2005. Diet of Barn Owl (*Tyto Alba*) in the Largest Subtropical Wetlands of Argentina and Eastern Paraguay. *Journal of Raptor Research* 39(1): 65-69.
- Pautasso, A.A. 2008 Mamíferos de la provincia de Santa Fe, Argentina. *Comunicaciones del Museo Provincial de Ciencias Naturales Florentino Ameghino* 13: 1-248.
- Romano, M., R. Biasitti and L. De Santis. 2002. Dieta de *Tyto alba* en una localidad urbana y otra rural en la Región Pampeana Argentina. *Hornero* 17(1): 25-29.
- Voglino, D., U.F.J. Pardiñas and P. Teta. 2005. *Holochilus chacarius chacarius* (Rodentia, Cricetidae) en la provincia de Buenos Aires, Argentina. *Mastozoología Neotropical* 11(2): 243-247.

RECEIVED: January 2010
 REVISED: June 2010
 ACCEPTED: August 2010
 PUBLISHED ONLINE: October 2010
 EDITORIAL RESPONSIBILITY: Ana Paula Carmignotto